### SRM VALLIAMMAI ENGINEERING COLLEGE

(An Autonomous Institution)

ESTD. 1999 - Approved by AICTE - Accredited by NBA
'A' Grade Accreditation by NAAC - Affiliated to Anna University
ISO 9001:2015 Certified Institution

# DEPARTMENT OF MATHEMATICS QUESTION BANK



#### I SEMESTER

M.E. Structural Engineering

1918103 - Advanced Mathematical Methods

**Regulation – 2019** 

Academic Year 2022- 2023

Prepared by

Mr. N. Sundarakannan, Assistant Professor/ Mathematics

Mr. L. Mohan, Assistant Professor/ Mathematics

### SRM VALLIAMMAI ENGNIEERING COLLEGE

(An Autonomous Institution)





SRM Nagar, Kattankulathur – 603203.

#### **DEPARTMENT OF MATHEMATICS**

Year & Semester : I/I

**Section** : M.E. (Structural Engineering)

**Subject Code** : 1918103

Subject Name : ADVANCED MATHEMATICAL METHODS

**Degree & Branch** : M.E – Structural Engineering

Staff in charge : Mr. L. Mohan

d				
	S.No	QUESTIONS	COMPETENCE	LEVEL

## UNIT -1 LAPLACE TRANSFORM TECHNIQUES FOR PARTIAL DIFFERENTIAL EQUATIONS

Laplace transform, Definitions, properties – Transform error function, Bessel's function, Dirac Delta function, Unit Step functions – Convolution theorem – Inverse Laplace Transform: Complex inversion formula – Solutions to partial differential equations: Heat equation. Wave equation.

	equation, Wave equation.			
1	PART A - 2 MARK QUESTIONS			
1.	Define Laplace Transform of a function	Remembering	BTL1	
2.	State the existence condition of Laplace Transform	Analyzing	BTL4	
3.	If $L(f(t) = F(s))$ , then prove that $L[e^{at} f(t)] = F(s-a)$	Applying	BTL3	
4.	Find $L[t\cos^2 t]$	Analyzing	BTL4	
5.	Determine $L\{2t-3e^{2t}\}$	Analyzing	BTL4	
6.	Find $L[e^{-4t} \sin 3t]$	Analyzing	BTL4	
7.	State Initial and Final value theorems for Laplace transform	Remembering	BTL1	
8.	$\operatorname{Find} L \left[ \frac{e^{-bt} - e^{-at}}{t} \right]$	Analyzing	BTL4	
9.	State and prove that change of scale property in Laplace transforms	Applying	BTL3	
10.	Define Laplace transform of Error function	Remembering	BTL1	
11.	Find Laplace transform of $t e^{-2t} \cos t$	Analyzing	BTL4	
12.	Define Bessel's function.	Applying	BTL3	
13.	Find $L^{-1}\left(\frac{s-5}{s^2+25}\right)$	Analyzing	BTL4	
14.	Show that $\int_0^t J_0(u)J_0(t-u)du = \sin t$	Understand	BTL2	
15.	Find the inverse Laplace transform of $\log \left( \frac{s+a}{s+b} \right)$	Analyzing	BTL4	
16.	Define unit step function and find its Laplace transform	Remembering	BTL1	

17.	Find the inverse Laplace transform of $\log \left( \frac{s+1}{s-1} \right)$	Applying	BTL3
18.	State Convolution theorem	Applying	BTL3
19.	Define Dirac- Delta function.	Applying	BTL3
20.	Find $L^{-1}\left(\frac{1}{(s+1)^2}\right)$	Analyzing	BTL4
21.	Find $L\{e^{-2t}\}$	Applying	BTL3
22.	Find $e^{-2t}\cos t$	Applying	BTL3
23.	Find $L(t \cos t)$	Applying	BTL3
24.	Find $L^{-1}\left(\frac{1}{(s-2)^2}\right)$	Applying	BTL3
25.	Find $\log\left(\frac{s+a}{s-b}\right)$	Applying	BTL3
	PART B - 13 MARK QUESTIONS		
1	Find $L[t \sin 5t \cos 2t]$	Applying	BTL3
2	Find the Laplace Transform of $erf(t^{\frac{1}{2}})$	Analyzing	BTL4
3	Find $L\left(\frac{\cos\sqrt{t}}{\sqrt{t}}\right)$	Evaluating	BTL5
4	Using Convolution theorem, find the inverse Laplace Transform of $\frac{s}{(s^2+9)(s^2+25)}$	Analyzing	BTL4
5	Find the Laplace transform of (i) $J_0(t)$ (ii) $tJ_0(t)$	Evaluating	BTL5
6	Using Convolution theorem, evaluating $L^{-1}(\frac{s}{(s^{2}+a)(s^{2}+1)})$	Applying	BTL3
7	Using Laplace transform, solve the IBVP $PDE: u_t = 3u_{xx},  BCs: u(\frac{\pi}{2}, t) = 0,  u_x(0, t) = 0,$ $ICs: u(x, 0) = 30\cos 5x.$	Evaluating	BTL5
8	Using Laplace transform, solve the IBVP $u_{tt} = u_{xx} \ 0 < x < l, t > 0, \ u(x,0) = 0, u_{t}(x,0) = \sin(\frac{\pi x}{l}), 0 < x < l,$ $u(0,t) = 0 \ and \ u(l,t) = 0, t > 0$	Evaluating	BTL5
9	Using Laplace transform, solve the IBVP $PDE: u_{tt} = u_{xx} \ 0 < x < 1, t > 0, \ BCs: u(x,0) = u(1,t) = 0, t > 0$ $ICs: u(x,0) = \sin \pi x, u_t(x,0) = -\sin \pi x, 0 < x < 1$	Evaluating	BTL5
10	Using Laplace transform solve the following IBVP $PDE: ku_t = u_{xx}$ , $0 < x < l$ , $0 < t < \infty$ , $BCS: u(0,t) = 0$ , $u(l,t) = g(t)$ , $0 < t < \infty$ , IC: u(x,0) = 0, $0 < x < l$ .	Analyzing	BTL4
11	Using Laplace transform method solve the initial boundary value problem described by	Evaluating	BTL5

$ \begin{array}{c} conditions \ u(0,t)=0 \ , u(x,0)=0, \\ u_r(0,t)=l \ for \ 0 \le x < \infty \ and \ u \ is \ bouded \ as \ x \to \infty. \end{array} $ $ \begin{array}{c} Applying \ the \ convolution \ theorem \ to \ evaluating \\ L^{-1} \left[ \begin{array}{c} s \\ (s^2+a^2)^2 \end{array}; t \right] \\ \\ 13  Derive \ the \ Laplace \ transform \ of \ Bessel \ functions \\ (i) \ J_0(x) \ (ii) \ J_1(x) \\ \\ 14  Find \ the \ inverse \ Laplace \ Transform \ of \ \frac{\sinh(x\sqrt{s})}{\sinh(t\sqrt{s})} \\ \text{in} \ 0 < x < l \ . \\ \\ 15  Find \ L[te^{-t}\cos 2t] \\ \\ \\ 16  Find \ the \ Laplace \ Transform \ of \ f(t) = \left\{ \begin{array}{c} \frac{t}{b}, \ for \ 0 < t < b \\ \frac{2b-t}{b}, \ for \ b < t < 2b \\ \frac{2b-t}{b}, \ for \ b < t < 2b \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ $		$u_{tt} = c^2 u_{xx}$ , $0 \le x < \infty$ , $t \ge 0$ , subject to the				
Applying the convolution theorem to evaluating $L^{-1}\left[\frac{s}{(s^2+a^2)^2};t\right]$ Analyzing BTL4  13 Derive the Laplace transform of Bessel functions $(i) J_0(x) (ii) J_1(x)$ Analyzing BTL4  14 Find the inverse Laplace Transform of $\frac{\sinh(x\sqrt{s})}{\sinh(t\sqrt{s})}$ Evaluating BTL5 in $0 < x < t$ .  15 Find $L^{-1}(t) = \frac{t}{t} \int_{t}^{t} for \ 0 < t < t$ Analyzing BTL3  16 Find the Laplace Transform of $f(t) = \frac{t}{t} \int_{t}^{t} for \ 0 < t < t$ Analyzing BTL3  17 Find $L^{-1}\left(\log\left(\frac{s^2-1}{s^2}\right)\right)$ Evaluating BTL5 Evaluating BTL5  18 Using Convolution theorem, find the inverse Laplace Transform of $\frac{s}{(s^2+16)(s^2+25)}$ Analyzing BTL4  19 Part C (15 Marks)  1. Verify Initial Value Theorem and Final Value Theorem for $f(t) = 1 + e^{-t} (\sin t + \cos t)$ Obtain the solution of free vibration of a semi infinite string governed by PDE $u_m = e^2 u_m \ 0 < x < \infty$ , $e > 0$ ICs: Creating BTL6  2. Find the inverse Laplace Transform of $\frac{1}{(s+1)(s-2)^2}$ by Analyzing BTL4		conditions $u(0,t) = 0$ , $u(x,0) = 0$ ,				
12 $L^{-1}\left[\frac{s}{(s^2+a^2)^2};t\right]$ Analyzing BTL4  13 Derive the Laplace transform of Bessel functions (i) $J_0(x)$ (ii) $J_1(x)$ Analyzing BTL4  14 Find the inverse Laplace Transform of $\frac{\sinh(x\sqrt{s})}{\sinh(l\sqrt{s})}$ Evaluating BTL5  15 Find $L[t e^{-t} \cos 2t]$ Applying BTL3  16 Find the Laplace Transform of $f(t) = \begin{cases} \frac{t}{b}, & for \ 0 < t < b \\ \frac{2b-t}{b}, & for \ b < t < 2b \end{cases}$ Analyzing BTL4  17 Find $L^{-1}\left(\log\left(\frac{s^2-1}{s^2}\right)\right)$ Evaluating BTL5  18 Using Convolution theorem, find the inverse Laplace Transform of $\frac{s}{(s^2+16)(s^2+25)}$ Analyzing BTL4  19 Part C (15 Marks)  1. Verify Initial Value Theorem and Final Value Theorem for $f(t) = 1 + e^{-t} (\sin t + \cos t)$ Obtain the solution of free vibration of a semi infinite string governed by PDE $u_n = c^2 u_n \ 0 < x < \infty$ , $c > 0$ ICs: $u(x,0) = f(x)$ , $u_t(x,0) = g(x)$ , $0 < x < \infty$ .  3. Find the inverse Laplace Transform of $\frac{1}{(s+1)(s-2)^2}$ by Analyzing BTL4		$u_t(0,t) = l \text{ for } 0 \le x < \infty \text{ and } u \text{ is bouded as } x \to \infty.$				
Derive the Laplace transform of Bessel functions $(i) \ J_0(x) \ (ii) \ J_1(x)$ Analyzing BTL4  14 Find the inverse Laplace Transform of $\frac{\sinh(x\sqrt{s})}{\sinh(l\sqrt{s})}$ Evaluating BTL5  15 Find $L[te^{-t}\cos 2t]$ Applying BTL3  16 Find the Laplace Transform of $f(t) = \begin{cases} \frac{t}{b}, \ for \ 0 < t < b \\ \frac{2b-t}{b}, \ for \ b < t < 2b \end{cases}$ Analyzing BTL4  17 Find $L^{-1}\left(\log\left(\frac{s^2-1}{s^2}\right)\right)$ Evaluating BTL5  18 Using Convolution theorem, find the inverse Laplace Transform of $\frac{s}{(s^2+16)(s^2+25)}$ Part C (15 Marks)  1. Verify Initial Value Theorem and Final Value Theorem for $f(t) = 1 + e^{-t}(\sin t + \cos t)$ Obtain the solution of free vibration of a semi infinite string governed by PDE $u_n = c^2 u_{xn} 0 < x < \infty$ , $c > 0$ ICs: $u(x,0) = f(x)$ , $u_x(x,0) = g(x)$ , $0 < x < \infty$ .  3. Find the inverse Laplace Transform of $\frac{1}{(s+1)(s-2)^2}$ by Analyzing BTL4		Applying the convolution theorem to evaluating				
Derive the Laplace transform of Bessel functions (i) $J_0(x)$ (ii) $J_1(x)$ BTL4  14 Find the inverse Laplace Transform of $\frac{\sinh(x\sqrt{s})}{\sinh(t\sqrt{s})}$ Evaluating BTL5 in $0 < x < t$ .  15 Find $L[te^{-t}\cos 2t]$ Applying BTL3  16 Find the Laplace Transform of $f(t) = \begin{cases} \frac{t}{b}, & for \ 0 < t < b \\ \frac{2b-t}{b}, & for \ b < t < 2b \end{cases}$ Analyzing BTL4 given that $f(t+2b)=f(t)$ for $t>0$ 17 Find $L^{-1}\left(\log\left(\frac{s^2-1}{s^2}\right)\right)$ Evaluating BTL5 Using Convolution theorem, find the inverse Laplace Transform of $\frac{s}{(s^2+16)(s^2+25)}$ Analyzing BTL4  18 Part C (15 Marks)  1. Verify Initial Value Theorem and Final Value Theorem for $f(t) = 1 + e^{-t}(\sin t + \cos t)$ Evaluating BTL5 Obtain the solution of free vibration of a semi infinite string governed by PDE $u_n = c^2 u_x, 0 < x < \infty$ , $c > 0$ ICs: Creating BTL6 $u(x,0) = f(x), \ u_t(x,0) = g(x), \ 0 < x < \infty$ .	12	$I^{-1} \left[ \begin{array}{c} S \\ \end{array} \cdot t \right]$	Analyzing	BTL4		
Find the inverse Laplace Transform of $\frac{\sinh(x\sqrt{s})}{\sinh(t\sqrt{s})}$ Evaluating BTL5 in $0 < x < l$ .  15 Find $L[te^{-t}\cos 2t]$ Applying BTL3  16 Find the Laplace Transform of $f(t) = \begin{cases} \frac{t}{b}, & \text{for } 0 < t < b \\ \frac{2b-t}{b}, & \text{for } b < t < 2b \end{cases}$ Analyzing BTL4  17 Find $L^{-1}\left(\log\left(\frac{s^2-1}{s^2}\right)\right)$ Evaluating BTL5  18 Using Convolution theorem, find the inverse Laplace Transform of $\frac{s}{(s^2+16)(s^2+25)}$ Analyzing BTL4  19 Part C (15 Marks)  10 Verify Initial Value Theorem and Final Value Theorem for $f(t) = 1 + e^{-t} (\sin t + \cos t)$ Obtain the solution of free vibration of a semi infinite string governed by PDE $u_n = c^2 u_\infty 0 < x < \infty$ , $c > 0$ ICs: Creating BTL6  2. Find the inverse Laplace Transform of $\frac{1}{(s+1)(s-2)^2}$ by Analyzing BTL4						
Find the inverse Laplace Transform of $\frac{\sinh(x\sqrt{s})}{\sinh(l\sqrt{s})}$ Evaluating BTL5 in $0 < x < l$ .  15 Find $L[te^{-t}\cos 2t]$ Applying BTL3  16 Find the Laplace Transform of $f(t) = \begin{cases} \frac{t}{b}, & for \ 0 < t < b \\ \frac{2b-t}{b}, & for \ b < t < 2b \end{cases}$ Analyzing BTL4 given that $f(t+2b)=f(t)$ for $t>0$ 17 Find $L^{-1}\left(\log\left(\frac{s^2-1}{s^2}\right)\right)$ Evaluating BTL5  18 Using Convolution theorem, find the inverse Laplace Transform of $\frac{s}{(s^2+16)(s^2+25)}$ Analyzing BTL4  19 Part C (15 Marks)  1. Verify Initial Value Theorem and Final Value Theorem for $f(t) = 1 + e^{-t} (\sin t + \cos t)$ Obtain the solution of free vibration of a semi infinite string governed by PDE $u_n = c^2 u_{xx} 0 < x < \infty, c > 0$ ICs: $u(x,0) = f(x), u_t(x,0) = g(x), 0 < x < \infty$ .  3. Find the inverse Laplace Transform of $\frac{1}{(s+1)(s-2)^2}$ by Analyzing BTL4	13		Analyzing	BTI 4		
	13	$(i) \ J_0(x) \ (ii) \ J_1(x)$		DIE!		
		Find the inverse Laplace Transform of $\frac{\sinh(x\sqrt{s})}{\sqrt{s}}$				
15 Find $L[te^{-t}\cos 2t]$ Applying BTL3  16 Find the Laplace Transform of $f(t) = \begin{cases} \frac{t}{b}, & for \ 0 < t < b \\ \frac{2b-t}{b}, & for \ b < t < 2b \end{cases}$ Analyzing BTL4  given that $f(t+2b)=f(t)$ for $t>0$ 17 Find $L^{-1}\left(\log\left(\frac{s^2-1}{s^2}\right)\right)$ Evaluating BTL5  18 Using Convolution theorem, find the inverse Laplace Transform of $\frac{s}{(s^2+16)(s^2+25)}$ Analyzing BTL4  19 Part C (15 Marks)  1. Verify Initial Value Theorem and Final Value Theorem for $f(t) = 1 + e^{-t} (\sin t + \cos t)$ Evaluating BTL5  2. Obtain the solution of free vibration of a semi infinite string governed by PDE $u_n = c^2 u_{xx} 0 < x < \infty$ , $c > 0$ ICs: Creating BTL6  2. Find the inverse Laplace Transform of $\frac{1}{(s+1)(s-2)^2}$ by Analyzing BTL4	14		Evaluating	BTL5		
Find the Laplace Transform of $f(t) = \begin{cases} \frac{t}{b}, & for \ 0 < t < b \\ \frac{2b-t}{b}, & for \ b < t < 2b \end{cases}$ Analyzing BTL4 given that $f(t+2b)=f(t)$ for $t>0$ 17 Find $L^{-1}\left(\log\left(\frac{s^2-1}{s^2}\right)\right)$ Evaluating BTL5  18 Using Convolution theorem, find the inverse Laplace Transform of $\frac{s}{(s^2+16)(s^2+25)}$ Analyzing BTL4  19 Part C (15 Marks)  1. Verify Initial Value Theorem and Final Value Theorem for $f(t) = 1 + e^{-t} (\sin t + \cos t)$ Evaluating BTL5  2. String governed by PDE $u_{tt} = c^2 u_{xx} \ 0 < x < \infty, c > 0$ ICs: Creating BTL6  2. Find the inverse Laplace Transform of $\frac{1}{(s+1)(s-2)^2}$ by Analyzing BTL4		in 0 < x < l .				
given that $f(t+2b)=f(t)$ for $t>0$ 17 $Find L^{-1}\left(\log\left(\frac{s^2-1}{s^2}\right)\right)$ Evaluating BTL5  18 Using Convolution theorem, find the inverse Laplace Transform of $\frac{s}{(s^2+16)(s^2+25)}$ Analyzing BTL4  Part C (15 Marks)  1. Verify Initial Value Theorem and Final Value Theorem for $f(t)=1+e^{-t}(\sin t+\cos t)$ Evaluating BTL5  Obtain the solution of free vibration of a semi infinite string governed by PDE $u_{tt}=c^2u_{xx}0< x<\infty$ , $c>0$ ICs: Creating BTL6  1. Find the inverse Laplace Transform of $\frac{1}{(s+1)(s-2)^2}$ by Analyzing BTL4	15	Find $L[t e^{-t} \cos 2t]$	Applying	BTL3		
given that $f(t+2b)=f(t)$ for $t>0$ 17 $Find L^{-1}\left(\log\left(\frac{s^2-1}{s^2}\right)\right)$ Evaluating BTL5  18 Using Convolution theorem, find the inverse Laplace Transform of $\frac{s}{(s^2+16)(s^2+25)}$ Analyzing BTL4  1. Verify Initial Value Theorem and Final Value Theorem for $f(t)=1+e^{-t}(\sin t+\cos t)$ Evaluating BTL5  2. Obtain the solution of free vibration of a semi infinite string governed by PDE $u_{tt}=c^2u_{xx}0< x<\infty$ , $c>0$ ICs: Creating BTL6  2. Find the inverse Laplace Transform of $\frac{1}{(s+1)(s-2)^2}$ by Analyzing BTL4		$\left(\frac{t}{-}  for \ 0 < t < h\right)$				
given that $f(t+2b)=f(t)$ for $t>0$ 17 $Find L^{-1}\left(\log\left(\frac{s^2-1}{s^2}\right)\right)$ Evaluating BTL5  18 Using Convolution theorem, find the inverse Laplace Transform of $\frac{s}{(s^2+16)(s^2+25)}$ Analyzing BTL4  1. Verify Initial Value Theorem and Final Value Theorem for $f(t)=1+e^{-t}(\sin t+\cos t)$ Evaluating BTL5  2. Obtain the solution of free vibration of a semi infinite string governed by PDE $u_{tt}=c^2u_{xx}0< x<\infty$ , $c>0$ ICs: Creating BTL6  2. Find the inverse Laplace Transform of $\frac{1}{(s+1)(s-2)^2}$ by Analyzing BTL4	1.0	Find the Laplace Transform of $f(t) = \begin{cases} b, & \text{if } t \in \mathbb{R} \\ b, & \text{otherwise} \end{cases}$		D.T. 4		
given that $f(t+2b)=f(t)$ for $t>0$ 17 $Find L^{-1}\left(\log\left(\frac{s^2-1}{s^2}\right)\right)$ Evaluating BTL5  18 Using Convolution theorem, find the inverse Laplace Transform of $\frac{s}{(s^2+16)(s^2+25)}$ Analyzing BTL4  1. Verify Initial Value Theorem and Final Value Theorem for $f(t)=1+e^{-t}(\sin t+\cos t)$ Evaluating BTL5  2. Obtain the solution of free vibration of a semi infinite string governed by PDE $u_{tt}=c^2u_{xx}0< x<\infty$ , $c>0$ ICs: Creating BTL6  2. Find the inverse Laplace Transform of $\frac{1}{(s+1)(s-2)^2}$ by Analyzing BTL4	16	$\left  \frac{2b-t}{b} \right $ , for $b < t < 2b$	Analyzing	BTL4		
17Find $L^{-1}\left(\log\left(\frac{s^2-1}{s^2}\right)\right)$ EvaluatingBTL518Using Convolution theorem, find the inverse Laplace Transform of $\frac{s}{(s^2+16)(s^2+25)}$ AnalyzingBTL4Part C (15 Marks)1.Verify Initial Value Theorem and Final Value Theorem for $f(t) = 1 + e^{-t} (\sin t + \cos t)$ EvaluatingBTL52.Obtain the solution of free vibration of a semi infinite string governed by PDE $u_n = c^2 u_{xx} 0 < x < \infty$ , $c > 0$ ICs: $u(x,0) = f(x)$ , $u_t(x,0) = g(x)$ , $0 < x < \infty$ .CreatingBTL63.Find the inverse Laplace Transform of $\frac{1}{(s+1)(s-2)^2}$ byAnalyzingBTL4						
Using Convolution theorem, find the inverse Laplace $Transform of \frac{s}{(s^2 + 16)(s^2 + 25)}$ Part C (15 Marks)  1. Verify Initial Value Theorem and Final Value Theorem for $f(t) = 1 + e^{-t} (\sin t + \cos t)$ Dotain the solution of free vibration of a semi infinite string governed by PDE $u_u = c^2 u_{xx} 0 < x < \infty$ , $c > 0$ ICs: $u(x,0) = f(x),  u_t(x,0) = g(x),  0 < x < \infty.$ BTL6  3. Find the inverse Laplace Transform of $\frac{1}{(s+1)(s-2)^2}$ by Analyzing BTL4						
Transform of $\frac{s}{(s^2+16)(s^2+25)}$ Part C (15 Marks)  1. Verify Initial Value Theorem and Final Value Theorem for $f(t) = 1 + e^{-t} (\sin t + \cos t)$ Dobtain the solution of free vibration of a semi infinite string governed by PDE $u_{tt} = c^2 u_{xx} 0 < x < \infty$ , $c > 0$ ICs: $u(x,0) = f(x)$ , $u_t(x,0) = g(x)$ , $0 < x < \infty$ .  The semi-part of $\frac{1}{(s+1)(s-2)^2}$ by Analyzing BTL4  Analyzing BTL4	17	Find $L^{-1} \left( \log \left( \frac{s^2}{s^2} \right) \right)$	Evaluating	BTL5		
Transform of $\frac{s}{(s^2+16)(s^2+25)}$ Part C (15 Marks)  1. Verify Initial Value Theorem and Final Value Theorem for $f(t) = 1 + e^{-t} (\sin t + \cos t)$ Dobtain the solution of free vibration of a semi infinite string governed by PDE $u_{tt} = c^2 u_{xx} 0 < x < \infty$ , $c > 0$ ICs: $u(x,0) = f(x)$ , $u_t(x,0) = g(x)$ , $0 < x < \infty$ .  The semi-part of $\frac{1}{(s+1)(s-2)^2}$ by Analyzing BTL4  Analyzing BTL4		Using Convolution theorem, find the inverse Laplace				
Part C (15 Marks)  1. Verify Initial Value Theorem and Final Value Theorem for $f(t) = 1 + e^{-t} (\sin t + \cos t)$ 2. Obtain the solution of free vibration of a semi infinite string governed by PDE $u_{tt} = c^2 u_{xx} 0 < x < \infty$ , $c > 0$ ICs: $u(x,0) = f(x),  u_t(x,0) = g(x),  0 < x < \infty.$ Creating BTL6 $u(x,0) = f(x),  u_t(x,0) = g(x),  0 < x < \infty.$ 3. Find the inverse Laplace Transform of $\frac{1}{(s+1)(s-2)^2}$ by Analyzing BTL4	18	_	Analyzing	BTL4		
1. Verify Initial Value Theorem and Final Value Theorem for $f(t) = 1 + e^{-t} (\sin t + \cos t)$ Obtain the solution of free vibration of a semi infinite string governed by PDE $u_{tt} = c^2 u_{xx} 0 < x < \infty$ , $c > 0$ ICs: $u(x,0) = f(x)$ , $u_t(x,0) = g(x)$ , $0 < x < \infty$ .  Find the inverse Laplace Transform of $\frac{1}{(s+1)(s-2)^2}$ by  Analyzing BTL4		$(s^2 + 16)(s^2 + 25)$				
1. for $f(t) = 1 + e^{-t} (\sin t + \cos t)$ Evaluating  Obtain the solution of free vibration of a semi infinite  2. string governed by PDE $u_{tt} = c^2 u_{xx} 0 < x < \infty$ , $c > 0$ ICs: $u(x,0) = f(x)$ , $u_t(x,0) = g(x)$ , $0 < x < \infty$ .  BTL6  3. Find the inverse Laplace Transform of $\frac{1}{(s+1)(s-2)^2}$ by  Analyzing  BTL5  BTL6		Part C (15 Marks)				
Obtain the solution of free vibration of a semi infinite  2. string governed by PDE $u_{tt} = c^2 u_{xx} 0 < x < \infty$ , $c > 0$ ICs: $u(x,0) = f(x),  u_{tt}(x,0) = g(x),  0 < x < \infty.$ BTL6  3. Find the inverse Laplace Transform of $\frac{1}{(s+1)(s-2)^2}$ by  Analyzing  BTL4	1	<del>-</del>	Evaluating	BTL5		
2. string governed by PDE $u_{tt} = c^2 u_{xx} 0 < x < \infty$ , $c > 0$ ICs: Creating BTL6 $u(x,0) = f(x), \ u_t(x,0) = g(x), \ 0 < x < \infty.$ 3. Find the inverse Laplace Transform of $\frac{1}{(s+1)(s-2)^2}$ by Analyzing BTL4	1.		Dvaraating	BILS		
$u(x,0) = f(x),  u_t(x,0) = g(x),  0 < x < \infty.$ 3. Find the inverse Laplace Transform of $\frac{1}{(s+1)(s-2)^2}$ by Analyzing BTL4			~ .			
3. Find the inverse Laplace Transform of $\frac{1}{(s+1)(s-2)^2}$ by Analyzing BTL4	2.		Creating	BTL6		
(		$u(x,0) = f(x), u_t(x,0) = g(x), 0 < x < \infty.$				
(	3	Find the inverse Laplace Transform of $\frac{1}{(s+1)(s-2)^2}$ by	Analyzina	RTI /		
	3.		Anaryzing	DILA		
Solve the following IBVP using Laplace Transform		1				
PDE: $u_{x} = u_{xx}$ , $0 < x < 1$ , $t > 0$ :	4			DITT 2		
4. BCs: $u(0,t) = 1$ , $u(1,t) = 1$ , $t > 0$ Applying BTL3	4.	BCs: $u(0,t) = 1$ , $u(1,t) = 1$ , $t > 0$	Applying	B1L3		
ICs: $u(x,0) = 1 + \sin \pi x$ , $0 < x < 1$ .		ICs: $u(x,0) = 1 + \sin \pi x$ , $0 < x < 1$ .				
Solve the initial boundary value problem described by						
$u_{tt} = c^2 u_{xx}, \ 0 \le x < \infty, \ t \ge 0$		$u_{tt} = c^2 u_{xx}, \ 0 \le x < \infty, \ t \ge 0$				
5. subject to the conditions Evaluating BTL5	5	subject to the conditions	Evaluating	BTI 5		
$u(0,t) = 0 , u(x,0) = 0, u_t(0,t) = k  for  0 \le x < \infty$	] 3.	ŭ	Lvanuatilig	DILJ		
and u is bounded as $x \to \infty$ by using Laplace transform		and <i>u</i> is bounded as $x \to \infty$ by using Laplace transform				
method		method				

## UNIT -2 FOURIER TRANSFORM TECHNIQUES FOR PARTIAL DIFFERENTIAL EQUATIONS

Fourier transform: Definitions, properties – Transform of elementary functions, Dirac Delta function – Convolution theorem – Parseval's identity – Solutions to partial differential equations: Heat equation, Wave equation, Laplace and Poison's equations.

PART A - 2 MARK QUESTIONS				
1 Define Fourier transform of a function	Remembering	BTL1		
2 Find the Fourier transform of $f(x) = \begin{cases} \cos x, & \text{in } 0 < x < \pi \\ 0, & \text{otherwise} \end{cases}$	Remembering	BTL1		
3 State Fourier transform pairs	Analyzing	BTL4		
4 State and prove shifting property in Fourier transform	Applying	BTL3		
5 Find the Fourier transform of $e^{-a x }$ , $-\infty < x < \infty$	Remembering	BTL1		
Find the Fourier Transform of $f(x) = \begin{cases} 1, &  x  < a \\ 0, &  x  > a \end{cases}$	Analyzing	BTL4		
7 State and prove modulation property in Fourier transform	Applying	BTL3		
8 State and prove change of scale property in Fourier transform	Applying	BTL3		
9 State Fourier Cosine transform pair	Remembering	BTL1		
10 State Fourier Sine transform pair	Remembering	BTL1		
11 Find the Fourier Sine transform of $e^{-ax}$ , $0 < x < \infty$	Analyzing	BTL4		
12 Find the Fourier Sine transform of 1/x	Analyzing	BTL4		
13 Find the Fourier cosine transform of $e^{-x}$ , $0 < x < \infty$	Remembering	BTL1		
Determine the frequency spectrum of the signal $g(t) = f(t) \cos \omega_c t$	Applying	BTL3		
If $F(\alpha)$ is the Fourier Transform of $f(x)$ , then the Fourier Transform of $f(ax)$ is $\frac{1}{ a }F(\alpha/a)$	Applying	BTL3		
16 Find the Fourier transform of Dirac Delta function	Analyzing	BTL4		
17 State Convolution Theorem on Fourier Transform	Remembering	BTL1		
18 State Parseval's identity on Fourier Transform	Remembering	BTL1		
Find the Fourier cosine Transform of $f(x) = \begin{cases} x, & 0 < x < a \\ 0, & x > a \end{cases}$	Analyzing	BTL4		
If F(s) is the Fourier transform of $f(x)$ , then find the Fourier transform of $f(x)$ cos ax	Evaluating	BTL5		
21 Prove that $F_S[f(ax)] = \frac{1}{a}F_S\left(\frac{s}{a}\right)$	Understanding	BTL 2		
Find the Fourier cosine transform of $e^{-2x}$ , $0 < x < \infty$	Remembering	BTL1		
Find the Fourier Sine transform of $e^{-3x}$ , $0 < x < \infty$	Remembering	BTL1		
24 If $F(s) = F[f(x)]$ , then find $F[xf(x)]$ .	Understanding	BTL 2		
State and Prove any one Modulation theorem on Fourier Transform	Understanding	BTL 2		
, i	PART B – 13 MARK QUESTIONS			

1	Find the Fourier Transform of $f(x) = \begin{cases} 1 -  x , & \text{if }  x  < 1 \\ 0, & \text{if }  x  > 1 \end{cases}$ hence deduce that $\int_{0}^{\infty} \left[ \frac{\sin t}{t} \right]^{2} dt = \frac{\pi}{2}$	Analyzing	BTL4
2	Find the Fourier Cosine and Sine Transform of $e^{-2x}$ and evaluate the integrals $\int_{0}^{\infty} \frac{\cos \alpha x}{\alpha^2 + 4} d\alpha$	Analyzing	BTL4
3	Find the Fourier Transform of $f(x)$ defined by $f(x) = \begin{cases} 1 - x^2, & \text{if }  x  < 1 \\ 0, & \text{if }  x  > 1 \end{cases}$ Hence $\text{evaluate } \int_{0}^{\infty} \left( \frac{x \cos x - \sin x}{x^3} \right) \cos \frac{x}{2} dx.$	Analyzing	BTL4
4	Find the Fourier Transform of $f(x) = \begin{cases} 1, & \text{if }  x  < a \\ 0, & \text{if }  x  > a \end{cases}$ hence deduce that $\int_{0}^{\infty} \left[ \frac{\sin t}{t} \right] dt = \frac{\pi}{2} \text{ and } \int_{0}^{\infty} \left[ \frac{\sin t}{t} \right]^{2} dt = \frac{\pi}{2}$	Evaluating	BTL5
5	Using Fourier Sine Transform, prove that $\int_{0}^{\infty} \frac{x^2 dx}{(x^2 + 9)(x^2 + 16)} = \frac{\pi}{14}$	Evaluating	BTL5
6	A one-dimensional infinite solid, $-\infty < x < \infty$ , is initially at temperature F(x). For times t>0, heat is generated within the solid at a rate of g(x ,t) units, Determines the temperature in the solid for t>0	Evaluating	BTL5
7	A uniform string of length L is stretched tightly between two fixed points at $x=0$ and $x=L$ . If it is displaced a small distance $\in$ at a point $x=b$ , $0 < b < L$ , and released from rest at time $t=0$ , find an expression for the displacement at subsequent times.	Evaluating	BTL5
8	Solve the heat conduction problem described by $\frac{\partial u}{\partial t} = k \frac{\partial^2 u}{\partial x^2}, 0 < x < \infty, t > 0,  u(0,t) = u_0, t > 0$ $u(x,0) = 0 < x < \infty, u \text{ and } \frac{\partial u}{\partial x} \text{ both tend to zero as } x \to \infty$	Analyzing	BLT4
9	Find the steady state temperature distribution $u(x,y)$ in a long square bar of side $\pi$ with one face maintained at constant temperature $u_0$ and other faces at zero temperature.	Analyzing	BLT4
10	Using the method of integral transform, solve the following potential problem in the semi-infinite Strip	Analyzing	BTL4

1.	Using the Fourier Cosine Transform of $e^{-ax}$ & $e^{-bx}$ , show	Evaluating	BTL5
Part C - 15 Marks			
18	Find the Fourier Transform of $f(x)$ defined by $f(x) = \begin{cases} 1 - x^2, & \text{if }  x  < 1 \\ 0, & \text{if }  x  > 1 \end{cases} \text{ hence evaluate } \int_0^\infty \left( \frac{x \cos x - \sin x}{x^3} \right) dx.$	Evaluating	BTL5
17	Find the Fourier transform of $e^{-a x }$ if $a > 0$ , deduce that $\int_{0}^{\infty} \frac{dx}{(x^2 + a^2)} = \frac{\pi}{4a^3}$ , if $a > 0$	Evaluating	BTL5
16	Find the Fourier Cosine and Sine Transform of $e^{-2x}$ and evaluate the integrals $\int_{0}^{\infty} \frac{\alpha \sin \alpha x}{\alpha^2 + 4} d\alpha$	Analyzing	BTL4
15	Find the Fourier Transform of $f(x) = \begin{cases} 1 -  x , & \text{if }  x  < 1 \\ 0, & \text{if }  x  > 1 \end{cases}$ hence deduce that $\int_{0}^{\infty} (\frac{\sin t}{t})^{4} dt = \frac{\pi}{3}$	Analyzing	BTL4
14	Solve the following boundary value problem in the half-plane $y > 0$ described by PDE : $u_{xx} + u_{yy} = 0$ , $-\infty < x < \infty$ , $y > 0$ BCs : $u(x,0) = f(x)$ , $-\infty < x < \infty$ u is bounded as $y \to \infty$ ; $u$ and $\frac{\partial u}{\partial x}$ both vanish as $ x  \to \infty$	Evaluating	BTL5
13	Compute the displacement $u(x,t)$ of an infinite string using the method of Fourier transform given that the string is initially at rest and that the initial displacement is $f(x)$ , $-\infty < x < \infty$ .	Creating	BTL6
12	Solve the heat conduction problem described $\frac{\partial u}{\partial t} = k \frac{\partial^2 u}{\partial x^2}, \ 0 < x < \infty, t > 0 \ u(0,t) = u_0 \ t > 0,$ $u(x,0) = 0, \ 0 < x < \infty, \ u \ and \ \frac{\partial u}{\partial x}$ Both tends to $0 \ x \to \infty$	Analyzing	BTL4
11	Using the finite Fourier transform, solve the BVP described by PDE: $\frac{\partial V}{\partial t} = \frac{\partial^2 V}{\partial x^2}, 0 < x < 6, t > 0, \text{ subject to BC:}$ $V_x(0,t) = 0 = V_x(6,t), \text{ IC : } V(x,0) = 2x.$	Analyzing	BTL4
	described by $PDE: u_{xx} + u_{yy} = 0$ , $0 < x < \infty$ , $0 < y < a$ Subject to BCs: $u(x,0)=f(x)$ , $u(x,a)=0$ , $u(x,y)=0$ , $0 < y < a$ , $0 < x < \infty$ and $\frac{\partial u}{\partial x}$ tend to zero as $x \to \infty$		

	that $\int_0^\infty \frac{d\alpha}{(a^2 + \alpha^2)(b^2 + \alpha^2)} = \frac{\pi}{2ab(a+b)}$ . $a > 0, b > 0$		
2.	Determine the temperature distribution in the semi- infinite medium $x \ge 0$ . When the end $x = 0$ is maintained at zero temperature and the initial temperature distribution is $f(x)$ .	Creating	BTL6
3.	A infinitely long string having one end at $x=0$ is initially at rest along x-axis. The end $x=0$ is given transverse displacement $f(t)$ , when $t>0$ . Find the displacement of any point of the string at any time.	Evaluating	BTL5
4.	Solve the following Neumann Problem  PDE: $u_{xx}(x, y) + u_{yy}(x, y) = 0, -\infty < x < \infty$ BC : $u_y(x,0) = f(x), -\infty < x < \infty$ u and $\frac{\partial u}{\partial x}$ both vanish as $ x  \to \infty$ .	Creating	BTL6
5.	A uniform string of length $l$ is stretched tightly between two fixed points at $x=0$ and $x=l$ . If it is displaced a small distance $\in$ at a point $x=a$ , $0 < a < l$ , and released from rest at time $t=0$ , find an expression for the displacement at subsequent times.	Evaluating	BTL5

### **UNIT -3 CALCULUS OF VARIATIONS**

Concept of variation and its properties – Euler's equation – Functional dependant on first and higher order derivatives – Functionals dependant on functions of several independent variables – Variational problems with moving boundaries – Problems with constraints – Direct methods – Ritz and Kantorovich methods.

	PART A - 2 MARK QUESTIONS		
1	Define a functional	Remembering	BTL1
2	Write Euler's equation for functional.	Applying	BTL3
3	Write Euler-Poisson equation.	Remembering	BTL1
4	Find the exteremals of the functional $\int (y^2 - y'^2 - 2y \sin x) dx.$	Remembering	BTL1
5	Test for an extremum the functional Find the extremals of $\int_{x_0}^{x_1} (y^2 + y^{1^2} - 2ye^x) dx$	Applying	BTL3
6	Prove that the commutative character for the operators of variation and differentiation.	Applying	BTL3
7	Show that the symbols $d/dx$ and $\delta$ are commutative.	Remembering	BTL1
8	State Brachistochrone problem.	Remembering	BTL1
9	Find the curves on which the functional $\int_{0}^{1} (y'^{2} + 12xy) dx \text{ with } y(0) = 0 \text{ and } y(1) = 1 \text{ can be extremised.}$	Applying	BTL3
10	Write the ostrogradsky equation for the functional	Analyzing	BTL4

	$I[z(x,y)] = \iint_D \left\{ \left( \frac{\partial z}{\partial x} \right)^2 + \left( \frac{\partial z}{\partial y} \right)^2 + z f(x,y) \right\} dx dy$		
	Write the ostrogradsky equation for the functional		
11	$\iint_{D} \left( \left[ \frac{\partial z}{\partial x} \right]^{2} + \left[ \frac{\partial z}{\partial y} \right]^{2} \right) dx dy$	Analyzing	BTL4
12	Define isoperimetric problems.	Applying	BTL3
13	Define moving boundaries.	Remembering	BTL1
14	Find the transversality condition for the functional $v = \int_{x_0}^{x_1} A(x, y) \cdot \sqrt{1 + (\frac{dy}{dx})^2} dx$ with the right boundary moving along $y_1 = \varphi(x_1)$	Applying	BTL3
15	What is the differential equation the functional $I[y(x)] = \int_{0}^{1} (xy + y^{2} - 2y^{2}y') dx  y(0) = 1,  y(1) = 2.$	Evaluating	BTL5
16	Write down biharmonic equation.	Applying	BTL3
17	State Hamilton's Principle	Remembering	BTL1
18	Find the fundmental lemma of calculus of variation.	Remembering	BTL1
19	Explain direct methods in variational problem	Understand	BTL2
20	Explain Rayleigh-Ritz method.	Understand	BTL2
21	Find the euler equation of the functional $\int_0^{\pi/2} \left( y^2 - \left( \frac{dy}{dx} \right)^2 - 2y \sin x \right) dx$	Applying	BTL3
22	Determine the differential equation of the functional $\int_0^{\pi} (y')^2 - y^2 + 4y \cos x  dx$	Remembering	BTL1
23	What is the shortest distance between two points in a plane	Remembering	BTL1
24	Find extremal of the functional $I(y(x)) = \int_0^1 (1 + y''^2) dx$	Understand	BTL2
25	The transversality condition for the functional $w = \int_{x_0}^{x_1} G(x, y) \cdot \sqrt{1 + (y')^2} dx$ with the right boundary moving along $y_1 = f(x_1)$	Understand	BTL2
PART B - 13 MARK QUESTIONS			
1	$I[y(x)] = \int_{0}^{1} (xy + y^{2} - 2y^{2}y') dx , y(0) = 1, y(1) = 2.$	Analyzing	BTL4
2	Find the extremal of the function $v[y(x), z(x)] \int_{x_0}^{x} \left[2yz - 2y^2 + y^{'2} - z^{'2}\right] dx.$	Applying	BTL3
3	Show that the straight line is the sharpest distance between two points in a plane.	Analyzing	BTL4
4	Show that the curve which extremize $I = \int_0^{\frac{\pi}{2}} [(y^{11})^2 - y^2 + x^2] dx \text{ given that } y(0) = 0, y^1(0) = 1, y\left(\frac{\pi}{4}\right) = \frac{1}{\sqrt{2}}, y^1\left(\frac{\pi}{4}\right) = \frac{1}{\sqrt{2}} \text{ is } y = \sin x.$	Analyzing	BTL4

5	Find the shortest distance between the point (-1,5) and the parabola $y^2 = x$	Analyzing	BTL4
6	Find the extremals of the functional $\int_{0}^{1} (1 + y''^{2}) dx$ .	Evaluating	BTL5
	Subject to $y(0) = 0$ , $y'(0) = 1$ , $y(1) = 1$ , $y'(1) = 1$ .		
7	Find the extremal of the functional $\int_{0}^{\pi} (y^{2} - y^{2} + 4y \cos x) dx.  y(0) = 0  , y(\pi) = 0.$	Evaluating	BTL5
8	Solve the boundary value problem $y^{11} - y + x = 0 \ (0 \le x \le 1) \ y(0) = y(1) = 0$ by Rayleigh Ritz method.	Applying	BTL3
9	To determine the shape of a solid of revolution moving in a flow of gas with least resistance.	Analyzing	BTL4
10	Derive the Euler's equation and use it, find the extremals of the functional $V[y(x)] = \int_{x_0}^{x_1} (y^2 + y'^2 - 2y \sin x) dx$ .	Creating	BTL6
11	Find an approximate solution to the problem of the minimum, of the functional $J(y) = \int_{0}^{1} (y'^2 - y^2 + 2xy) dx$ . $y(0)=0=y(1)$ by Ritz method and compare it with the exact solution.	Evaluating	BTL5
12	Find the solid of maximum volume formed by the revolution of a given surface area.	Evaluating	BTL5
13	Find the extremals of the functional $v[y(x), z(x)] = \int_{0}^{\pi/2} (y'^2 + z'^2 + 2yz) dx$ . Given that $y(0) = 0$ , $y(\pi/2) = -1$ , $z(0) = 0$ , $z(\pi/2) = 1$	Evaluating	BTL5
14	Find the curve connect in the points A and B which is traversed by a particle sliding from A to B in the shortest time.	Creating	BTL6
15	Find the extremals of $\int_{x_0}^{x_1} (y^{1^2} + 2yy^1 - 16y^2) dx$	Evaluating	BTL5
16	Solve the extremal $v[y(x)] = \int_{x_0}^{x_1} \frac{dy}{dx} \left(1 + x^2 \frac{dy}{dx}\right) dx$ .	Evaluating	BTL5
17	Find the plane curve of a fixed perimeter and maximum area.	Applying	BTL3
18	Find the extremal of the functional $I(y,z) = \int_{0}^{1} (y^{2} + z^{2} + 2y) dx; \ y(0) = 1, \ y(1) = \frac{3}{2},$ $z(0) = 0, z(1) = 1)$	Analyzing	BTL4
	Part C – 15 Marks		
1.	Discuss the Branchistochrone Problem.	Remembering	BTL1
2.	Deterine the extermal of the functional	Applying	BTL3

	$I(y(x)) = \int_{-a}^{a} \left(\frac{1}{2}\mu y''^2 + \rho y\right) dx$ that satisfies the condition $y(-a) = 0, \ y'(-a) = 0, \ y(a) = 0, \ y'(a) = 0$		
3.	Prove that the sphere is the solid figure of revolution which for a given surface area has maximum volume.	Remembering	BTL1
4.	Using Ritz method, find the approximate solution of the problem of the minimum of the functional $I(y(x)) = \int_0^2 (y'^2 + y^2 + 2xy) dx , y(0) = y(2) = 0$ and compare with the exact solution.	Applying	BTL3
5	A curve C joining the points $(x_1, y_1)$ and $(x_2, y_2)$ is revolved about the x-axis. Find the shape of the curve C, so that the surface area generated is a minimum.	Remembering	BTL1

### UNIT-4 CONFORMAL MAPPING AND APPLICATIONS

Introduction to conformal mappings and bilinear transformations — Schwarz Christoffel transformation — Transformation of boundaries in parametric form — Physical applications : Fluid flow and heat flow problems.

TTUIC	now and near now problems.		
	PART A - 2 MARK QUESTIONS		
1	Define Conformal Mapping.	Remembering	BTL1
2	Define Isogonal transformation.	Remembering	BTL1
3	Find the stagnation points of the flow represented by the complex potential $\Omega(z) = z^2$	Remembering	BTL1
4	Define Mobius transformation	Remembering	BTL1
5	Define cross ratio of $Z_1$ , $Z_2$ , $Z_3$ , $Z_4$	Understand	BTL2
6	Define Schwarz-Christoffel transformation.	Applying	BTL3
7	Define invariant point of the transformation $w = f(z)$	Remembering	BTL1
8	Find the fixed points of the transformation $w = \frac{z-2}{z+4}$	Applying	BTL3
9	Find the critical point of the transformation $W^2 = (Z - \alpha)(Z - \beta)$	Analyzing	BTL4
10	Find the fixed points of $w = \frac{3z+1}{z-1}$	Analyzing	BTL4
11	Let C be a curve in the z-plane with parametric equations $x=F(t)$ , $y=G(t)$ . Show that the transformation $z=F(\omega)+i$ $G(\omega)$ maps the curve C on to the real axis of the w-plane.	Evaluating	BTL5
12	Discuss the transformation defined by w= coshz.	Remembering	BTL1
13	Define equipotential lines.	Remembering	BTL1
14	State the formula for Schwarz Christoffel transformation	Understand	BTL2
15	Define Heat flux of heat flow.	Remembering	BTL1
16	State the basic assumptions of fluid flow.	Remembering	BTL1
17	Define streamline	Remembering	BTL1
18	Show that for a closed polygon, the sum of exponents in Schwarz Christoffel transformation is -2.	Applying	BTL3

19	Define complex temperature of heat flow	Understand	BTL2
20	Define velocity potential, source and sink	Remembering	BTL1
21	Find the critical point of the transformation $w = z + \frac{1}{z}$	Remembering	BTL1
22	Find the invariant points of $w = \frac{1+z}{1-z}$	Remembering	BTL1
23	What is the sum of the exterior angles of any closed polygon?	Understand	BTL2
24	Define Irrotational flow.	Remembering	BTL1
25	When we will say the fluid is incompressible?	Remembering	BTL1
	PART B - 13 MARK QUESTIONS		
1	Define Bilinear transformation and find the bilinear	Understanding	BTL2
	transformation which maps $z_1 = -2$ , $z_2 = 0$ , $z_3 = 2$ on to the points $w_1 = \infty$ , $w_2 = 1/2$ , $w_3 = 3/4$ .		
2	Find a transformation which maps a polygon in the w-plane to the unit circle in the z-plane.	Applying	BTL3
	$Q$ $-\pi+i\pi$ $R$ $Q'$ $R'$ $x$ $y$		
3	Find a BLT $z_1 = 0$ , $z_2 = -i$ , $z = -1$ in to the point s $w_1 = 1$ , $w_2 = 1$ $w_3 = 0$	Remembering	BTL-1
4	Find the bilinear transformation which maps the points z=1, i, -1 into the points w=2, i, -2. Find the fixed and critical points of the transformation.	Understand	BTL2
5	Prove that a harmonic function $\Phi(x, y)$ remains harmonic under the transformation $w=f(z)$ where $f(z)$ is analytic and $f'(z) \neq 0$ .	Understand	BTL2
6	Find the transformation which maps the semi-infinite strip bounded by v=0, u=0 and v=b into the upper half of the z-plane.	Analyzing	BTL4
7	Find the complex potential due to a source at $z = -a$ and a sink at $z = a$ equal strengths $k$ , Determine the equipotential lines and streamlines and represent graphically and find the speed of the fluid at any point.	Evaluating	BTL5
8	Discuss the motion of a fluid having complex potential $\Omega(z) = ik \log z$ where $k > 0$	Applying	BTL3
	The complex potential of a fluid flow is given by	Evaluating	BTL5

9	$\Omega(z) = V_0(z + \frac{a^2}{z})$ where $V_0$ and $\alpha$ are positive constants.		
	<ul> <li>(a) Obtain equations for the streamlines and equipotential lines, represent them graphically and interpret physically.</li> <li>(b) Show that we can interpret the flow as that around a circular obstacle of radius a.</li> <li>(c) Find the velocity at any point and determine its value far</li> </ul>		
	from the obstacle		
10	(d) Find the stagnation points.		7.77
10	Fluid emanates at a constant rate from an infinite line source perpendicular to the z plane at z=0 (a) Show that the speed of the fluid at a distance r from the source is V=k/r, where k is a constant	Analyzing	BTL4
	(b) Show that the complex potential is $\Omega(z) = k \ln z$ . (c) What modification should be made in (b) if the line source is at z=a?		
	(d) What modification is made in (b) if the source is replaced by a sink in which fluid is disappearing at a constant rate?		
11	Find the complex potential for a fluid moving with constant speed $V_0$ in a direction making an angle $\delta$ with the positive x-axis. Also determine the velocity potential and stream function.	Applying	BTL3
12	Find the transformation which transforms the semi infinite strip bounded by $v=0$ , $v=\pi$ and $u=0$ onto the upper half zplane.	Analyzing	BTL4
13	Show that the transformation $w = \frac{3-z}{z-2}$ transforms circle with center(5/2, 0) and radius 1/2 in the z-plane into the	Applying	BTL3
	imaginary axis in w-plane and the interior of the circle into the right half of the plane.		
14	Find the transformation that maps an infinite strip $0 \le v \le \pi$ of the w-plane onto the upper half of z-plane.	Analyzing	BTL4
15	Find a bilinear transformation which maps points $z=0$ , -i, -1 into $w=i$ , 1, 0 respectively.	Applying	BTL3
16	Find the function harmonic in the upper half of the z- plane, which takes the prescribed values on x- axis given by $G(x) = \begin{cases} 1, & x > 0 \\ 0, & x < 0 \end{cases}$	Analyzing	BTL4
17	Prove that the bilinear transformation transforms circles of the z plane into circles of the w plane, where by circles we include circles of infinite radius which are straight lines.	Applying	BTL3
18	The bilinear transformation of the points which maps $0, 1,\infty$ on to -5, -1, -3	Analyzing	BTL4

	Part C- 15 Marks			
1.	Find the complex potential for a fluid moving with constant speed $V_0$ in a direction making an angle $\delta$ with the positive x axis. Also (i) Determine the velocity potential and stream function (ii)Determine the equations for the streamlines and equipotential lines.	Evaluating	BTL5	
2.	Discuss the Schwarz- Christoffel transformation.	Analyzing	BTL4	
3.	Find the Bilinear transformation which maps the points $z=1$ , i, -1 onto $w=i$ , 0, -i. Hence find the image of $ z <1$ and the invariant points of the transformation.	Applying	BTL3	
4.	Discuss the different types of Fluid flow and Heat Flow.	Analyzing	BTL4	
5.	Determine the transformation of the following figure  B  A  A  C  D  A  B'  C'  D'	Applying	BTL3	
	z - plane w - plane			

### **UNIT-5 TENSOR ANALYSIS**

14

Summation convention – Contravariant and covaraiant vectors – Contraction of tensors – Innerproduct – Quotient law – Metric tensor – Chrirstoffel symbols – Covariant differentiation – Gradient, divergence and curl.

**PART A - 2 MARK QUESTIONS: Define Tensor and Summation convention** Remembering BTL1 Define Kronecker delta function of tensor BTL1 Remembering 3 Define Covariant and Contravariant vectors Remembering BTL1 Define contraction of a tensor. 4 **Applying** BTL3 is a contravariant vector, 5 If  $A_i$  is a covariant and  $B^i$ **Applying** BTL3 show that  $A_i B^i$  is an invariant 6 State the Quotient law of tensors. BTL3 **Applying** Define reciprocal tensors. 7 BTL1 Remembering 8 Remembering BTL1 Prove that  $[pr,q]+[qr,p]=\frac{\partial g_{pq}}{\partial r^r}$ . Define Symmetric and Skew-symmetric tensors Remembering BTL1 10 Define divergence of a contra variant vector Remembering BTL1 11 Prove that Kronecker delta ( $\delta$ ) is a mixed tensor of order 2. **Analyzing** BTL4 12 Define Metric tensor and Conjugate tensor BTL3 **Applying** 13 State the Christoffel symbol of first and second kind. Understand BTL2

**Applying** 

BTL3

If f is a function of 'n' variables  $x^i$  then write the differential

	of f using summation convention.		
15	Write the terms contained in $S = a_{ij}x^ix^j$ , taking n=3.	Applying	BTL3
16	If $(ds)^2 = (dr)^2 + r^2(d\theta)^2 + r^2\sin^2\theta(d\phi)^2$ , find the value of [22.1] and [13,3]	Creating	BTL6
17	Prove that $\operatorname{div} A_i = \frac{1}{\sqrt{g}} \frac{\partial (\sqrt{g} A^k)}{\partial x^k}$ .	Analyzing	BTL4
18	Define Curl of a contra variant vector.	Applying	BTL3
19	Given $A^i$ and $B^j$ are contra variant vectors, show that	Applying	BTL3
	$A^i B^j$ is contra variant tensor of rank 2		
20	Prove that the property of the Christoffel symbols $\Gamma_{ij,k} + \Gamma_{kj,i} = \frac{\partial g_{ik}}{\partial q^j}$	Analyzing	BTL4
21	If $\varphi$ is a function of the <i>n</i> quantities $x^i$ , Find the differential of $\varphi$ using summation convention.	Applying	BTL3
22	Find $\delta_j^i \cdot \delta_k^j$	Remembering	BTL1
23	What is the type of the velocity of a fluid at any point?	Remembering	BTL1
24	What is the matrix form of a metric tensor of a line element in spherical polar coordinates?	Remembering	BTL1
25	Find $[pr,q]+[qr,p]$	Remembering	BTL1
	PART B – 13 MARK QUESTIONS		
1	If $A^i$ is a contravariant vector and $B_j$ is a covariant vector,	Evaluating	BTL5
	prove that $A_i B^j$ is a mixed tensor of order 2.		
2	Find $g$ and $g^{ij}$ corresponding to the metric	Analyzing	BTL4
	$(ds)^{2} = 5(dx^{1})^{2} + 3(dx^{2})^{2} + 4(dx^{3})^{2} - 6dx^{1}dx^{2} + 4dx^{2}dx^{3}$		
3	Find the components of the first and second fundamental tensor in spherical coordinates.	Creating	BTL6
4	Find the components of the metric tensor and the conjugate tensor in the cylindrical co-ordinates.	Applying	BTL3
5	Define Christoffel's symbols. Show that they are not tensors	Analyzing	BTL4
6	A covariant tensor has components $x + y$ , $xy$ , $2z - y^2$ in rectangular co-ordinates. Find its covariant components in cylindrical co-ordinates.	Evaluating	BTL5
7	Given the covariant components in rectangular co-	Analyzing	BTL4
	ordinates $2x - z$ , $x^2y$ , $yz$ . Find the covariant components in Spherical polar co-ordinates $(r, \theta, \varphi)$		
8	Show that the tensors $g_{ij}$ , $g^{ij}$ and the Kroneckor delta $\delta_j^i$ are	Applying	BTL3
9	constants with respect to covariant differentiation. Find the <i>g</i> and $g^{ij}$ for the metric $ds^2 = 5(dx^1)^2 + 3(dx^2)^2 + 4(dx^3)^2 - 6dx^1dx^2 + 4dx^2dx^3 + 0dx^3dx^1.$	Evaluating	BTL5
10	If $(ds)^2 = (dr)^2 + r^2(d\theta)^2 + r^2 \sin^2 \theta (d\phi)^2$ , find the value of	Evaluating	BTL5

	(i) [22,1] and [13,3] (ii) $\begin{cases} 1 \\ 22 \end{cases}$ and $\begin{cases} 3 \\ 13 \end{cases}$ Determine the metric tensors $g_{ij}$ , $g^{ij}$ in cylindrical co-		
11	Determine the metric tensors $g_{ij}$ , $g^{ij}$ in explination $g_{ij}$	Evaluating	BTL5
	ordinates.		
12	Prove that the covariant derivative of $g^{ij}$ is zero.	Evaluating	BTL5
13		Evaluating	BTL5
	Prove that (i) $\frac{\partial g_{ij}}{\partial x^k} = [ik, j] + [jk, i]$	Dvardating	BILS
	( ) 63		
	(ii) $\frac{\partial g^{ij}}{\partial x^k} = -g^{jl} \begin{Bmatrix} i \\ lk \end{Bmatrix} - g^{im} \begin{Bmatrix} j \\ mk \end{Bmatrix}$		
14	Given the metric	Evaluating	BTL5
	$ds^{2} = (dx^{1})^{2} + (x^{1})^{2} (dx^{2})^{2} + (x^{1})^{2} \sin^{2}(x^{2}) (dx^{3})^{2}$		
	find the Christoffel symbols [22, 1], [32, 3]		
	$\begin{bmatrix} 2 \\ 2 \end{bmatrix} \text{ and } \begin{bmatrix} 3 \\ 3 \end{bmatrix}$		
15	A covariant tensor has components $xy$ , $2y-z^2$ , $xz$ in	Evaluating	BLT5
	rectangular coordinates. Find its covariant components in		
	spherical coordinates		
16	Prove that the covariant derivative of $g^{ij}$ for the metric	Analyzing	BTL4
	$ds^{2} = 5(dx^{1})^{2} + 3(dx^{2})^{2} + 4(dx^{3})^{2} - 6dx^{1}dx^{2} + 4dx^{2}dx^{3} + 0dx^{3}dx^{1} $ is		
	zero.		
17	State and prove Quotient law of tensor.	Creating	BTL6
18	Given the covariant components in rectangular co-	Applying	BTL3
	ordinates $2x - z$ , $x^2y$ , $yz$ . Find the covariant components in		
	Cylindrical co-ordinates ( $\rho, \varphi, z$ )		
	Part C – 15 Marks		
1.	Show that any innerpoduct of the tensor $A_m^l$ and $B_r^{pq}$ is a	Creating	BTL6
	tensor of rank three.		
2.	Explain short notes about metric tensor with suitable	Analyzing	BTL4
	example.		
3.	Discuss the Curl and Divergent of a tenor.	Analyzing	BTL4
4.	A covariant tensor has components xy, 2y-z², xz in	Applying	BTL3
	rectangular coordinates. Find its covariant components in		
	spherical coordinates.		
5.	If g denotes the determinant of $g_{ij}$ then prove that	Analyzing	BTL4
	$\begin{cases} i \\ i \end{cases} = \frac{\partial}{\partial x^i} (\log \sqrt{g})$		

\*\*\*\*\*\*\*\*